

## **EXECUTIVE SUMMARY**

### **Introduction**

This is a recovery plan (Plan) for the protection and restoration of Middle Columbia River steelhead (*Oncorhynchus mykiss*), which spawn and rear in tributaries to the Columbia River in central and eastern Washington and Oregon (Figure ES-1). The Middle Columbia River steelhead distinct population segment (DPS) was listed as threatened under the Endangered Species Act of 1973 (ESA) on January 5, 2006 (71 FR 834).

Section 4(f) of the ESA requires NOAA’s National Marine Fisheries Service (NMFS) to develop recovery plans for marine species listed under the Act. Recovery plans identify actions needed to restore threatened and endangered species to the point that they are again self-sustaining elements of their ecosystems and no longer need the protections of the ESA. Although recovery plans are guidance, not regulatory documents, the ESA clearly envisions recovery plans as the central organizing tool for guiding each species’ recovery process. Recovery planning is an opportunity to search for the common ground, to organize protection and restoration of salmonid habitat, and to secure the economic and cultural benefits that accrue to human communities from healthy watersheds and rivers.

Eighteen of the 33 salmon and steelhead species in the Northwest region are listed as threatened or endangered. The Middle Columbia steelhead is among those with the best prospects of recovery, although it will require considerable political will and investment of long-term effort and funding. Modeling of the potential effects of the actions that are proposed in this plan (see Chapter 9) predicts that the DPS can achieve a “negligible” risk of extinction within a reasonable time frame – e.g. 25 to 50 years – if the actions are taken and if they have the predicted effects on steelhead habitat and survival. Cautious though this statement may be, it is a beacon of hope in the complex realm of salmonid recovery in the Northwest. The following sections tell the story.

### **ESA Requirements**

ESA section 4(a)(1) lists factors for re-classification or delisting that are to be addressed in recovery plans:

- A. The present or threatened destruction, modification, or curtailment of [the species’] habitat or range
- B. Over-utilization for commercial, recreational, scientific or educational purposes
- C. Disease or predation
- D. The inadequacy of existing regulatory mechanisms
- E. Other natural or human-made factors affecting its continued existence




Figure ES-1. Geographic boundaries of the Middle Columbia River Steelhead DPS, showing land ownership.

ESA section 4(f)(1)(B) directs that recovery plans, to the extent practicable, incorporate:

1. a description of such site-specific management actions as may be necessary to achieve the plan's goal for the conservation and survival of the species;
2. objective, measurable criteria which, when met, would result in a determination, in accordance with the provisions of this chapter, that the species be removed from the list; and;
3. estimates of the time required and the cost to carry out those measures needed to achieve the plan's goal and to achieve intermediate steps toward that goal.

In addition, it is important for recovery plans to provide the public and decision makers with a clear understanding of the goals and strategies needed to recover a listed species and the science underlying these conclusions (NMFS Interim Recovery Planning Guidance, October 2004).

Once a species is deemed recovered and therefore removed from a listed status, section 4(g) of the ESA requires the monitoring of the species for a period of not less than 5 years to ensure that it retains its recovered status.

### **Steelhead Distribution and Life History**

The spawning range of the Middle Columbia steelhead DPS extends over an area of approximately 35,000 square miles in the Columbia plateau of eastern Washington and eastern Oregon. The DPS includes all naturally spawned populations of steelhead in streams from above the Wind River, Washington, and the Hood River, Oregon (exclusive), upstream to, and including, the Yakima River, Washington, excluding steelhead from the Snake River Basin (64 FR 14517; 71 FR 849). The Cascade Mountains form the western border of the plateau in both Oregon and Washington, while the Blue Mountains form the eastern edge. The southern border is marked by the divides that separate the upper Deschutes and John Day basins from the Oregon High Desert and drainages to the south. The Wenatchee Mountains and Palouse areas of eastern Washington border the Middle Columbia on the north.

Most of the region is privately owned (64 percent), with the remaining area under Federal (23 percent), tribal (10 percent) and state (3 percent) ownership (Figure ES-1). Most of the landscape consists of rangeland and timberland, with significant concentrations of dryland agriculture in parts of the range. Irrigated agriculture and urban development are generally concentrated in valley bottoms. Human populations in these regions are growing.

Four artificial propagation programs are considered part of the DPS: the Touchet River Endemic Summer Steelhead Program, the Yakima River Kelt Reconditioning Program, and the Umatilla River and Deschutes River steelhead hatchery programs.

The species *Oncorhynchus mykiss* exhibits perhaps the most complex suite of life history traits of any species of Pacific salmonid. These fish can be anadromous (migratory) or freshwater residents (and under some circumstances, apparently yield offspring of the opposite form). Steelhead can spawn more than once (iteroparous), whereas all other *Oncorhynchus* except cutthroat trout (*O. clarki*) spawn once and then die (semelparous).

Within the range of West Coast steelhead, spawning migrations occur throughout the year, with seasonal peaks of activity. The “runs” are usually named for the season in which the peak occurs. Most steelhead can be categorized as one of two run types, based on their sexual maturity when they re-enter freshwater and how far they go to spawn. In the Pacific Northwest, summer steelhead enter freshwater between May and October and require several months to mature before spawning; winter steelhead enter freshwater between November and April with well-developed gonads and spawn shortly thereafter. Summer steelhead usually spawn farther upstream than winter steelhead (Withler 1966; Roelofs 1983; Behnke 1992).

The Middle Columbia River steelhead DPS includes populations of inland winter steelhead in the Klickitat River, White Salmon River, Fifteenmile Creek, and possibly Rock Creek.

### **Relationship of Steelhead DPS to Resident *O. mykiss***

“Steelhead” is the name commonly applied to the anadromous (migratory) form of the biological species *Oncorhynchus mykiss*. The common name of the non-anadromous, or resident, form is rainbow trout. When NMFS originally listed the Middle Columbia River steelhead as threatened on March 25, 1999 (64 FR 14517), it was classified as an “evolutionarily significant unit” (ESU) of salmonids that included both the anadromous and resident forms. Recently, NMFS revised its species determinations for West Coast steelhead under the ESA, delineating anadromous, steelhead-only “distinct population segments” (DPS). NMFS listed the Middle Columbia River steelhead DPS as threatened on January 5, 2006 (71 FR 834). Rainbow trout are under the jurisdiction of the U.S. Fish and Wildlife Service (USFWS). This recovery plan addresses steelhead and not rainbow trout, as is consistent with the 2006 ESA listing decision.

### **Context of Plan Development**

While NMFS is directly responsible for ESA recovery planning for salmon and steelhead, NMFS believes that ESA recovery plans for salmon and steelhead should be based on the many state, regional, tribal, local, and private conservation efforts already underway throughout the region. Local support of recovery plans by those whose activities directly affect the listed species, and whose actions will be most affected by recovery efforts, is essential. NMFS therefore supports and participates in locally led collaborative efforts to develop recovery plans that involve local communities, state, tribal, and Federal entities, and other stakeholders.

This Plan is the product of a collaborative process initiated by NMFS with assistance from the Middle Columbia Forum (Mid-C Forum), a bi-state, tri-tribe group convened by NMFS to provide input on the development of the DPS recovery plan. NMFS developed this Plan by drawing upon the best available scientific information provided by the six regional recovery plans included as appendices to this Plan (i.e. the management unit plans, described below and in Section 1.6.), and by a regional team of scientists (the Interior Columbia Technical Recovery Team, described below). The draft plan went through repeated reviews and revisions in response to comments from both the scientific team and the Mid-C Forum. Participants in the Mid-C Forum include the Oregon Department of Fish and Wildlife (ODFW), Washington Department of Fish and Wildlife (WDFW), the Yakama Nation, Confederated Tribes of the Warm Springs Indian Reservation, Confederated Tribes of the Umatilla Indian Reservation, Washington Governor’s Salmon Recovery Office, Oregon Governor’s Natural Resources Office, Snake River Salmon Recovery Board, Yakima Basin Fish and Wildlife Recovery Board, US Bureau of

Reclamation (BOR), US Fish and Wildlife Service (USFWS), US Forest Service (USFS), US Army Corps of Engineers (COE), Klickitat County, and NMFS Northwest Region.

### **Tribal Trust and Treaty Responsibilities**

Northwest Indian Tribes have legally enforceable treaty rights reserving to them a share of the salmon harvest. A complex history of treaties, executive orders, legislation, and court decisions have culminated in the recognition of tribes as co-managers who share management responsibilities and rights for fisheries in the Columbia Basin.

Ensuring a sufficient abundance of salmon and steelhead to sustain harvest is an important element in fulfilling trust responsibilities and treaty rights as well as garnering public support for recovery plans. ESA and tribal trust responsibilities complement one another. Both depend on a steady upward trend toward ESA recovery and delisting in the near term, while making aquatic habitat, harvest, and land management improvements for the long-term.

### **Recovery Domains and Technical Recovery Teams**

Currently, there are 18 ESA-listed ESUs/DPSs of Pacific salmon and steelhead in the Pacific Northwest. For the purpose of recovery planning for these species, NMFS Northwest Region designated five geographically based “recovery domains” (Figure ES-2). The range of the Middle Columbia River steelhead DPS is located in the Middle Columbia sub-domain of the Interior Columbia domain.

For each domain, NMFS appointed a team of scientists, nominated for their geographic and species expertise, to provide a solid scientific foundation for recovery plans. The charge of each Technical Recovery Team (TRT) is to define ESU/DPS structures, develop recommendations on biological viability criteria for each ESU or DPS and its component populations, provide scientific support to local and regional recovery planning efforts, and provide scientific evaluations of proposed recovery plans. The Interior Columbia TRT (ICTRT) includes biologists from NMFS, states, and academic institutions.

### **Viable Salmonid Populations**

All the TRTs used the same biological principles for developing their recommendations for ESU/DPS and population viability criteria – criteria that may be used, along with criteria based on mitigation of the factors for decline, in determining whether a species has recovered sufficiently to be downlisted or delisted. These principles are described in a NMFS technical memorandum, *Viable Salmonid Populations and the Recovery of Evolutionarily Significant Units* (McElhany et al. 2000).

Viable salmonid populations (VSP) are defined in terms of four parameters: abundance, productivity or growth rate, spatial structure, and diversity. A viable ESU/DPS is naturally self-sustaining, with a high probability of persistence over a 100-year time period. Each TRT made recommendations using the VSP framework, based on data availability, the unique biological characteristics of the ESUs/DPSs and habitats in the domain, and the members’ collective




Figure ES-2. Columbia Basin Recovery Domains for NMFS Northwest Region

experience and expertise. Although NMFS has encouraged the TRTs to develop regionally specific approaches for evaluating viability and identifying factors limiting recovery, all the TRTs are working from a common scientific foundation. Viability criteria are an important part of recovery goals, as described later in this summary.

### Management Units

In each domain, NMFS worked with state, tribal, local, and other Federal entities to develop planning forums that build to the extent possible on ongoing, locally led recovery efforts. NMFS defined “management units” based on jurisdictional boundaries as well as areas where local planning efforts were underway (Figure ES-3). The Middle Columbia management units are (1) Oregon; (2) Washington Gorge, which, in turn, is subdivided into three planning areas, White Salmon, Klickitat, and Rock Creek; (3) Yakima subbasin; and (4) Southeast Washington.

### Management Unit Recovery Plans

Although NMFS has prepared this plan for the entire DPS, the DPS plan’s component parts (the management unit plans, Appendices A-F) are the work of local groups and county, state, Federal, and tribal entities within the Middle Columbia River region on both sides of the river.

- **Oregon Management Unit: *Conservation and Recovery Plan for Oregon Steelhead Populations in the Middle Columbia River Steelhead Distinct Population Segment (Oregon Steelhead Recovery Plan) (Appendix A).***

The Oregon Department of Fish and Wildlife (ODFW) is the lead for the Oregon Steelhead Recovery Plan. ODFW drew together three groups to help with the plan: the Middle Columbia Recovery Planning Team, made up of ODFW staff biologists and representatives from eight state natural resource agencies; a planning forum, the Middle Columbia Sounding Board, made up of representatives of local communities, agricultural water users, Federal and non-Federal land managers, governing bodies, tribes, and industry and environmental interests; and an Expert Panel of 12 biologists to examine limiting factors and threats for the 10 independent steelhead populations in Oregon.

- **Washington Gorge Management Unit: *Recovery Plans for the Klickitat (Appendix B), Rock Creek (Appendix C), and White Salmon (Appendix D) subbasin populations of Middle Columbia River steelhead***

Since there is not presently a Washington State sponsored salmon recovery planning board for this area, NMFS staff drafted the three plans in collaboration with the Yakama Nation, Washington Department of Fish and Wildlife, Klickitat County, the Washington State Governor's Salmon Recovery Office, other Federal agencies, state agencies, local governments, and the public.

- **Yakima Management Unit: *Yakima Steelhead Recovery Plan (Appendix E)***

The Yakima Basin Fish and Wildlife Recovery Board (YBFWRB), which includes representatives from the Yakama Nation, Benton, Kittitas, and Yakima counties, and 18 of the 24 municipalities in the Yakima Basin, developed the Yakima Basin Steelhead Recovery Plan (available at [www.ybfwrb.org](http://www.ybfwrb.org)).

- **Southeast Washington Management Unit: *Snake River Salmon Recovery Plan for Southeast Washington (Appendix F)***

The Snake River Salmon Recovery Board developed the Southeast Washington Recovery Plan. The Board consists of representatives of the Confederated Tribes of the Umatilla Indian Reservation; a county commissioner and citizen representative from Asotin, Columbia, Garfield, Walla Walla and Whitman counties; a land owner representative from Asotin, Columbia and Garfield counties; and the Walla Walla county irrigation district. The Board appointed a Regional Technical Team for technical and scientific assistance.




Figure ES-3. Management Units and Populations for the Middle Columbia River Steelhead DPS

## **Important Concepts in Steelhead (and Salmon) Biology**

Salmonid species' homing propensity (their tendency to return to the locations where they originated) creates unique patterns of genetic variation and connectivity that mirror the distribution of their spawning areas across the landscape. Diverse genetic, life history, and morphological characteristics have evolved over generations, creating runs highly adapted to diverse environments. It is this variation that gives the species as a whole the resilience to persist over time.

Historically, a salmon ESU or steelhead DPS typically contained multiple populations connected by some small degree of genetic exchange by straying spawners. Thus, the overall biological structure of the ESU/DPS is hierarchical; spawners in the same area of the same stream will share more characteristics than those in the next stream over. Fish whose natal streams are separated by hundreds of miles will have less genetic similarity.

### **Definition of Evolutionarily Significant Units/Distinct Population Segments**

An ESU or DPS is a distinctive group of Pacific salmon or steelhead that is uniquely adapted to a particular area or environment and cannot be replaced. Because of the hierarchical structure of salmonid populations, the concept of "distinctive group" has received considerable attention and refinement. An ESU is defined as a group of Pacific salmon that is "substantially reproductively isolated from other conspecific units and represents an important component of the evolutionary legacy of the species" (Waples 1991). A "population segment" is considered distinct (a DPS and hence a "species" for purposes of conservation under the ESA) if it is discrete from and significant to the remainder of its species based on factors such as physical, behavioral, or genetic characteristics; it occupies an unusual or unique ecological setting; or its loss would represent a significant gap in the species' range.

ESUs/DPSs may contain multiple populations that are connected by some degree of genetic exchange through "straying," and hence may have a broad geographic range across watersheds and river basins.

### **Major Population Groups**

Within an ESU/DPS, independent populations can be grouped into larger populations that share similar genetic, geographic, and/or habitat characteristics (McClure et al. 2003). These "major groupings" of populations (MPGs) are isolated from one another over a longer time scale than that defining the individual populations, but retain some degree of connectivity greater than that between ESUs/DPSs.

### **Independent Populations**

McElhany et al. (2000) defined an independent population as follows:

*"...a group of fish of the same species that spawns in a particular lake or stream (or portion thereof) at a particular season and which, to a substantial degree, does not interbreed with fish from any other group spawning in a different place or in the same place at a different season."*

## **Abundance and Productivity**

Abundance refers to spawners (adults on the spawning ground), measured over a time series, i.e. some number of years. The ICTRT often uses a recent 10- or 12-year geometric mean of natural spawners as a measure of current abundance.

The productivity of a population (the average number of surviving offspring per parent) is a measure of the population's ability to sustain itself. Productivity can be measured as spawner:spawner ratios (returns per spawner or recruits per spawner) (or adult progeny to parent), annual population growth rate, or trends in abundance. Population-specific estimates of abundance and productivity are derived from time series of annual estimates, typically subject to a high degree of annual variability and sampling-induced uncertainties.

Abundance and productivity are linked, as populations with low productivity can still persist if they are sufficiently large, and small populations can persist if they are sufficiently productive. A viable population needs sufficient abundance to maintain genetic health and to respond to normal environmental variation, and sufficient productivity to enable the population to quickly rebound from periods of poor ocean conditions or freshwater perturbations.

## **Spatial Structure and Diversity**

Spatial structure and diversity considerations are combined in the evaluation of a salmonid population's status because they often overlap. A population's spatial structure is made up of both the geographic distribution of individuals in the population and the processes that generate that distribution (McElhany et al. 2000, p. 18). Diversity refers to the distribution of traits within and among populations. Some traits are completely genetically based, while others, including nearly all morphological, behavioral, and life history traits, vary as a result of a combination of genetic and environmental factors (ibid., p. 19).

Populations with restricted distribution and few spawning areas are at a higher risk of extinction as a result of catastrophic environmental events, such as a landslide, than are populations with more widespread and complex spatial structures. Population-level diversity is similarly important for long-term persistence. Populations exhibiting greater diversity are generally more resilient to short-term and long-term environmental changes.

## **Middle Columbia Steelhead Populations and Major Population Groups**

The ICTRT (McClure et al. 2003) identified 20 historical populations of Middle Columbia steelhead, shown in Figure ES-4. This identification was based on genetic information, geography, life history traits, morphological traits, and population dynamics. Seventeen of these populations are extant, and three extirpated (White Salmon River, Deschutes Crooked River above Pelton Dam, and Willow Creek).

The ICTRT stratified the Middle Columbia River steelhead populations into MPGs based on ecoregion characteristics, life history types, and other geographic and genetic considerations. It identified four MPGs: Cascades Eastern Slope Tributaries, Yakima Basin, John Day Basin, and Umatilla/Walla Walla. The John Day Basin MPG is wholly within Oregon and the Yakima Basin

MPG is wholly within Washington. The other two include populations on both sides of the Oregon/Washington boundary.

### **Population Size**

Middle Columbia steelhead spawn in a wide range of tributary drainage areas, from small creeks, e.g. Fifteenmile Creek or Rock Creek, to very large rivers, such as the Lower John Day. The ICTRT categorized historical population sizes as Basic, Intermediate, Large, and Very Large, and set minimum abundance thresholds for viable steelhead populations of each type (ICTRT 2007a and 2007b). The abundance thresholds are associated with minimum productivity thresholds. Abundance and productivity are linked, within limits; above a certain threshold, higher productivity can compensate for lower abundance and vice versa. Table ES-1 shows the minimum abundance and productivity thresholds for the Middle Columbia steelhead populations to have a 95 percent probability of persistence for the next 100 years.

Proposed Middle Columbia Steelhead ESA Recovery Plan  
September 2008



Figure ES-4. Middle Columbia River Steelhead Populations and Major Population Groups.

**Table ES-1. Middle Columbia steelhead size categories (ICTRT 2007a)**

<b>Major Population Grouping</b>	<b>Population</b>	<b>Population Size</b>	<b>Abundance Threshold</b>	<b>Productivity Threshold</b>
Cascades Eastern Slope Tributaries	White Salmon (functionally extirp.)	Basic	500	1.56
	Klickitat R.	Intermediate	1000	1.35
	Fifteenmile Cr.	Basic	500	1.56
	Deschutes R. East	Intermediate	1000	1.35
	Deschutes R. West	Large <sup>1</sup>	1500	1.26
	Rock Cr.	Basic	500	1.56
	Crooked River (Extirp.)	Very Large	2250	1.19
	John Day River	Lower Mainstem JD	Very Large	2250
North Fork John Day		Large	1500	1.26
Middle Fork John Day		Intermediate	1000	1.35
South Fork John Day		Basic	500	1.56
Upper Mainstem JD		Intermediate	1000	1.35
Umatilla / Walla Walla Rivers	Umatilla R.	Large	1500	1.26
	Walla Walla R.	Intermediate	1000	1.35
	Touchet R.	Intermediate	1000	1.35
	Willow Crk. (Extirp.)	Intermediate	1000	1.35
Yakima River Group	Satus Cr.	Intermediate <sup>2</sup>	1000	1.35
	Toppenish Cr.	Basic	500	1.56
	Naches R.	Large	1500	1.26
	Upper Yakima	Large	1500	1.26

## Recovery Goals and Delisting Criteria

The recovery goals that are incorporated into a locally developed recovery plan may include delisting, reclassification (e.g., from endangered to threatened), and/or other “broad sense” goals that may go beyond the requirements for delisting to address, for example, other legislative mandates or social, economic, or ecological values. NMFS’ delisting criteria may include both technical and policy considerations. Delisting criteria must meet the ESA requirements, while recovery may be defined more broadly. A third “term of art” used in this recovery plan is recovery “scenarios” (Section 3.3). Recovery scenarios are combinations of viability status for individual populations within the DPS that will meet the ICTRT criteria for overall DPS viability.

Recovery criteria are of two kinds: the biological viability criteria, which deal with the VPS parameters at the population, MPG, and DPS levels, and the “threats” criteria, which relate to the five listing factors detailed in the ESA (see Sections 1.3 and 3.4 of this Plan). The threats criteria define the conditions under which the listing factors, or threats, can be considered to be

<sup>1</sup> This population is treated as Intermediate in size with respect to abundance and productivity criteria because of constraints on currently accessible habitat (i.e. Pelton Dam).

<sup>2</sup> For the historical population analysis, the ICTRT included the mainstem Yakima habitat below the confluence of Satus Creek in the Satus Creek population, making it Intermediate in size. However, if the mainstem component is lumped instead with mainstem Yakima River habitat upstream of Satus, the Satus Creek population would drop to Basic size. The Yakima Steelhead Recovery Plan discusses this question in more detail.

addressed or mitigated. Together these make up the “objective, measurable criteria” required under section 4(f)(1)(B) for the delisting decision.

The delisting criteria are based on the best available scientific information and incorporate the most current understanding of the DPS and the threats it faces. As this recovery plan is implemented, additional information will become available that can increase certainty about whether the threats have been abated, whether improvements in population and DPS status have occurred, and whether linkages between threats and changes in salmon status are understood. These criteria will be assessed through an adaptive management program under development for the Plan, and NMFS will thoroughly review the criteria during its 5- and 10-year status reviews of the DPS.

### **Biological Viability Criteria**

In 2007, the ICTRT completed its Technical Review Draft of *Viability Criteria for Application to Interior Columbia Basin Salmonid ESUs* (ICTRT 2007a). Biological viability criteria describe DPS characteristics associated with a low risk of extinction for the foreseeable future. These criteria are expressed in terms of the VSP parameters of abundance, productivity, spatial distribution, and diversity, according to guidelines developed by NOAA’s Northwest Fisheries Science Center and published as a NOAA Technical Memorandum, *Viable Salmonid Populations and the Recovery of Evolutionarily Significant Units* (McElhany et al. 2000; ICTRT 2007a). The ICTRT calculated varying levels of risk of extinction and related the risk levels to their criteria.

### **DPS Viability**

Since MPGs are geographically and genetically cohesive groups of populations, they are critical components of ESU or DPS spatial structure and diversity. Having all MPGs within a DPS at low risk provides the greatest probability of persistence for the DPS.

#### **DPS Viability Criterion (ICTRT 2007a)**

**All extant MPGs and any extirpated MPGs critical for proper functioning of the ESU/DPS<sup>3</sup> should be at low risk.**

### **MPG Viability**

MPG viability depends on the number, spatial arrangement, and diversity associated with its component populations.

---

<sup>3</sup> The Middle Columbia steelhead DPS has four extant and no extirpated MPGs. The three extirpated populations are addressed as part of the MPG-level criteria.

**MPG-Level Viability Criteria  
(ICTRT 2007a)**

The following five criteria should be met for an MPG to be regarded as at low risk (viable):

1. At least one-half of the populations historically within the MPG (with a minimum of two populations) should meet viability standards.
2. At least one population should be classified as “Highly Viable.”
3. Viable populations within an MPG should include some populations classified (based on historical intrinsic potential) as “Very Large,” “Large,” or “Intermediate,” generally reflecting the proportions historically present within the MPG. In particular, Very Large and Large populations should be at or above their composite historical fraction within each MPG.
4. All major life history strategies (e.g. spring and summer-run timing) that were present historically within the MPG should be represented in populations meeting viability requirements.
5. Remaining MPG populations should be maintained with sufficient abundance, productivity, spatial structure, and diversity to provide for ecological functions and to preserve options for ESU/DPS recovery.

The DPS criterion requiring viable populations in each of the extant MPGs would result in sustainable production across a substantial range of environmental conditions. The presence of viable populations across MPGs would preserve a high level of diversity within the DPS, thereby promoting long-term evolutionary potential for adaptation to changing conditions. The presence of multiple, relatively nearby, viable and maintained populations acts as protection against long-term impacts of localized catastrophic loss by serving as a source of re-colonization (ICTRT 2007a).

### **Population Viability**

To be viable, populations should meet criteria for all four VSP parameters (abundance, productivity, spatial structure, and diversity).

#### *Abundance and productivity*

The ICTRT defined abundance and productivity criteria for Middle Columbia steelhead populations (ICTRT 2005 and 2007) based on analyses of the intrinsic potential of the historically available habitat, the locations and sizes of major and minor spawning areas, and, within these areas, the abundance and productivity relationships that would result in a probability of low risk of extinction within 100 years (see Table ES-1 above). The abundance “thresholds” shown in the table represent the number of spawners needed for a population of the given size category to achieve the 5 percent (low) risk level at a given productivity.

#### *Spatial structure and diversity*

The spatial structure and diversity criteria are specific to each population, and are based on historical spatial distribution and diversity, to the extent these can be known or inferred. The

ICTRT cautions that there is a good deal of uncertainty in assessing the status of spatial structure and diversity in a population (ICTRT 2007a; McElhany et al. 2000).

### **Recovery Scenarios**

The risk levels of the populations within the DPS collectively determine MPG viability and, in turn, the likely persistence of the DPS. The ICTRT recommended that all MPGs in a DPS should be viable; however, it may not be necessary for all of the populations to attain the lowest risk level. There may be more than one way for a DPS to meet the viability criteria.

The ICTRT, in a January 8, 2007 technical memorandum (ICTRT 2007a), offered a detailed discussion of possible recovery scenarios for each MPG. They cautioned against closing off the options for any population prematurely, however, because of the many uncertainties in predicting the biological response to recovery actions. The ICTRT concluded that “a low risk strategy will target more populations than the minimum for viability” (ICTRT 2007a).

The management unit plans include locally determined recovery goals as well as viability criteria for the individual steelhead populations and MPGs in each management unit. Most of the plans also provide targets or objectives to measure progress within specified time frames, e.g. 10 to 50 years.

### **Threats Criteria**

At the time of a delisting decision for the Middle Columbia steelhead, NMFS will examine whether the section 4(a)(1) listing factors have been addressed. To assist in this examination, NMFS will use the listing factors (or threats) criteria described in Section 4.3 of this plan, in addition to evaluation of biological recovery criteria and other relevant data and policy considerations. It is possible that currently perceived threats will become insignificant in the future because of changes in the natural environment or changes in the way threats affect the entire life cycle of salmon. Consequently, NMFS expects that the relative priority of threats will change over time and that new threats may be identified. During the status reviews, NMFS will evaluate and review the listing factor criteria as they apply at that time. NMFS expects that if the proposed actions described in the Plan are implemented, they will make substantial progress toward meeting the listing factor (threats) criteria for the Middle Columbia steelhead.

### **Current Status Assessment**

The status of a salmonid ESU or DPS is expressed in terms of likelihood of persistence over 100 years, or in terms of risk of extinction within 100 years. The ICTRT defines viability at two levels: less than 5 percent risk of extinction within 100 years (viable) and less than 1 percent risk of extinction within 100 years (highly viable). A third category, “maintained,” represents a less than 25 percent risk. The risk level of the DPS is built up from the aggregate risk levels of the populations and MPGs. All four VSP parameters must be taken into account to determine the risk level.

Table ES-2 summarizes current status of the Middle Columbia steelhead populations, showing 10-year geometric mean abundance by population, estimated productivity, and the minimum abundance threshold needed for long-term viability. The table also includes the 10-year

geometric mean proportion of hatchery spawners for the populations where data are available, and the risk ratings of high, moderate, low, and very low, for abundance and productivity combined, and spatial structure and diversity combined. Figure ES-5 is a matrix combining all four parameters to illustrate the overall current risk rating of each population.

### **Current Population Status**

According to the ICTRT viability criteria, the majority of natural Middle Columbia steelhead populations are rated at moderate risk for all four VSP parameters – abundance, productivity, spatial structure, and diversity (Table ES-2 and Figure ES-5). This DPS includes one highly viable population (North Fork John Day), two viable (Fifteenmile Creek and Deschutes River Eastside), and three at high risk of extinction within 100 years (Deschutes Westside, Upper Yakima Mainstem, and Naches River).

### **MPG Status**

The viability ratings of the component populations of each Middle Columbia steelhead MPG are shown in Figure ES-5. None of the MPGs as a whole reaches low risk status according to the ICTRT's MPG-level criteria.

### **DPS Status**

The ICTRT's DPS-level viability criterion is that all extant MPGs and any extirpated MPGs critical for proper functioning of the DPS should be at low risk (ICTRT 2007). Thus, the Middle Columbia steelhead DPS does not currently meet viability criteria based on the determination that the four component MPGs are not at low risk.

*Proposed Middle Columbia Steelhead ESA Recovery Plan  
September 2008*

**Table ES-2. Middle Columbia River steelhead DPS populations: summary of abundance, productivity, risk ratings, and minimum abundance thresholds (Source: ICTRT 2007c and 2008). (Numbers subject to periodic updates as additional information becomes available.)**

<i>Population</i>	<i>Abundance Threshold<sup>4</sup></i>	<i>Size Category</i>	<i>Run Timing</i>	<i>10-year Geomean abundance</i>	<i>Abundance Range</i>	<i>10-yr Hatchery Fraction<sup>5</sup></i>	<i>Productivity<sup>6</sup></i>	<i>Productivity Standard Error</i>	<i>A&amp;P Risk<sup>7</sup> Rating</i>	<i>SSD Risk Rating</i>
<b>Eastern Cascades MPG</b>										
Deschutes (westside)	1000 <sup>8</sup>	Large (Inter)	Summer	456	108-1283	0.26	1.05	0.15	H	M
Deschutes (eastside)	1000	Intermed.	Summer	1599	299-8274	0.39	1.89	0.27	L	M
Klickitat River	1000	Intermed.	Wtr & Smr						M	M
Fifteenmile Creek	500	Basic	Winter	703	231-1922	0	1.82	0.20	L	L
Rock Creek	500	Basic	Summer	Insufficient Data					H	M
White Salmon	500	Basic		Functionally extirpated					N/A	N/A
Crooked River	2250	Very Large	Summer	Extirpated						
<b>Yakima River MPG</b>										
Upper Yakima River	1500	Large	Summer	85	34-283	0.02	1.12	0.22	H	H
Naches River	1500	Large	Summer	472	142-1454	0.06	1.12	0.22	H	M
Toppenish River	500	Basic	Summer	322	44-1252	0.06	1.60	0.30	M	M
Satus Creek (trib only)	1000	Intermed.	Summer	379	138-1000	0.06	1.73	0.14	M	M
<b>John Day Basin MPG</b>										
Lower Mainstem John Day	2250	Very Large	Summer	1800	563-6257	0.1	2.99	0.24	M	M
North Fork John Day	1500	Large	Summer	1740	369-10,235	0.08	2.41	0.22	VL	L
Upper Mainstem John Day	1000	Intermed.	Summer	524	185-5169	0.08	2.14	0.33	M	M
Middle Fork John Day	1000	Intermed.	Summer	756	195-3538	0.08	2.45	0.16	M	M
South Fork John Day	500	Basic	Summer	259	76-2729	0.08	2.06	0.27	M	M
<b>Umatilla/Walla Walla MPG</b>										
Umatilla River	1500	Large	Summer	1472	592-3542	0.36	1.50	0.15	M	M
Walla Walla Mainstem	1000	Intermed.	Summer	650	270-1746	0.02	1.34	0.12	M	M
Touchet River	1000	Intermed.	Summer	Insufficient Data					H	M
<b>Willow Creek</b>	1000	Intermed.	Summer	Extirpated					N/A	N/A


<sup>4</sup> Abundance threshold for viability based on habitat intrinsic potential

<sup>5</sup> Average proportion of hatchery spawners over most recent 10 years in the data series.

<sup>6</sup> Geomean return per spawner calculated over most recent 20 years in data series.

<sup>7</sup> H = high risk, M = moderate risk, L = low risk, VL = very low risk

<sup>8</sup> The Deschutes Westside steelhead population is classified as Large in terms of spatial structure, but its abundance threshold may be considered 1000 or 1500 because of “currently accessible area” considerations. See Carmichael, Richard W., Draft Recovery Plan for Oregon’s Middle Columbia River Steelhead: Progress Report. January 17, 2006



**Figure ES-5. Viability Ratings for Middle Columbia Steelhead Populations by MPG (developed by NMFS based on ICTRT 2008)**

### Gap Analysis

The ICTRT assessed the difference between a listed species’ or population’s current status for abundance and productivity and the viability criteria. This difference is called the “gap.” The gap, as used in this plan, is a measure, although it is inevitably imprecise, of the improvement in survival needed to meet viability criteria. As such, it is also an indicator of the level of effort needed to achieve recovery.

The ICTRT calculated the gap for each extant Middle Columbia steelhead population based on current abundance and productivity for the listed salmon and steelhead in the Interior Columbia Basin (ICTRT 2007c). They estimated the minimal survival rate changes needed for Middle Columbia steelhead populations to meet the abundance and productivity viability criteria for a 5 percent risk of extinction in a 100-year time frame.

In addition, the ICTRT (2007) estimated gaps under three different early-ocean survival scenarios; historical ocean conditions (ocean conditions that fish experienced over the past 60 years), pessimistic ocean conditions (ocean conditions experienced by the 1975-1997 brood years), and recent ocean conditions (ocean conditions experienced by fish during the 20-year assessment period). The ICTRT also estimated gaps assuming three different hydropower scenarios. However, only the base hydro condition, which assumed

that survival rates from the most recent 20 years would continue into the future, is reported here. (See NMFS 2008a for details on survival through the FCRPS under proposed improvements.)

A positive number, e.g. 21 percent gap for Eastern Cascades MPG, means the populations’ overall survival needs to increase 21 percent over current conditions to achieve viability criteria. A zero or negative number would mean there is no gap – the population currently meets viability criteria.

The analysis showed that none of the MPGs would be able to achieve a 5 percent or less risk of extinction over 100 years without recovery actions (Table ES-3). The Yakima Basin MPG shows the largest gap (77 percent) and also contains two historically large populations now at high risk of extinction, the Upper Yakima River and Naches River populations.

**Table ES-3. Median survival gap for the major population groups of the Middle Columbia Steelhead DPS (assuming recent ocean and base hydrosystem conditions and 5 percent risk)**

Eastern Cascades MPG	21 percent
John Day MPG	9 percent
Umatilla/Walla Walla MPG Sufficient data for only two of the three populations:	
Umatilla	9 percent
Walla Walla	34 percent
Yakima	77 percent

### **Limiting Factors and Threats**

The reasons for a species’ decline are generally described in terms of limiting factors and threats. NMFS defines limiting factors as the biological and physical conditions that limit a species’ viability – e.g., high water temperature – and defines threats as those human activities or natural processes that cause the limiting factors. For example, removing the vegetation along the banks of a stream can cause higher water temperatures, because the stream is no longer shaded. The threats contributing to the limiting factors and causes for a species’ decline are often described in terms of the “four Hs” – habitat (usually relating to the effects of land use and tributary water use), hydropower, harvest, and hatcheries. While the term “threats” carries a negative connotation, it does not mean that activities identified as threats are inherently undesirable. They are typically legitimate and necessary human activities that may at times have unintended negative consequences on fish populations—and that can also be managed in a manner that minimizes or eliminates the negative impacts.

Designing effective recovery strategies and actions requires understanding limiting factors and threats across the species’ entire life cycle and across the four Hs. This plan describes limiting factors and threats for the Middle Columbia steelhead DPS as a whole at a general level, and notes the most salient specific conditions that affect individual populations and limit the viability of specific MPGs. More detail is available in the individual management unit plans (Appendices A through F).

## **Limiting Factors and Threats for the DPS**

At a general level, based on information from the ICTRT and the four management unit plans, the major factors limiting the viability of Middle Columbia steelhead populations are *degraded tributary habitat, impaired fish passage in the mainstem Columbia River and tributaries, hatchery-related effects, and predation/competition/disease*. Two other factors, degradation of estuarine and nearshore marine habitat and harvest-related effects, pose some risk to steelhead viability for the entire DPS, but less than the other factors. Climate change represents a potentially significant threat to recovery of Middle Columbia steelhead populations (see ISAB 2007 and Section 6.3.8 of this plan).

## **Limiting Factors and Threats for the MPGs**

The MPG-level summaries of limiting factors are based on population-level summaries compiled from the relevant management unit plans.

### **Cascade Eastern Slope Tributaries MPG**

The following are major limiting factors for the Cascades Eastern Slope Tributaries MPG (see also the Oregon Steelhead Recovery Plan [Appendix A] and the Klickitat Recovery Plan [Appendix B]):

*Tributary habitat.* Degraded tributary habitat is a limiting factor to a greater or lesser degree throughout the area, including degraded riparian areas, reduced recruitment of large woody debris (LWD), altered sediment routing, low or altered stream flows, degraded water quality (especially high water temperatures), impaired floodplain connectivity/function, altered channel structure/complexity, and impaired fish passage.

*Mainstem passage.* Mainstem Columbia River hydro system effects are least for the Fifteenmile Creek and Klickitat River populations, which pass only one mainstem dam. The Deschutes River populations pass two mainstem dams, and the Rock Creek population passes three.

*Hatchery related effects.* Influence from hatchery fish could be a significant factor for this MPG because of out-of-subbasin straying onto natural spawning grounds in the Deschutes River and also because of potential effects of hatchery releases on naturally produced steelhead in the Klickitat River. The Oregon Mid-C Expert Panel considered out-of-subbasin (and out-of-DPS) hatchery strays a primary threat to genetic traits and productivity of naturally produced Deschutes river steelhead populations. Out-of-DPS hatchery strays comprised an estimated average of 29 percent of the Eastside population and 15.2 percent of the Westside population since 1990 (ICTRT 2008). This high fraction resulted in moderate risk ratings for spawner composition for both populations.

*Blocked migration to historically accessible habitat.* Historically, summer steelhead had free access to most of the Deschutes watershed. Currently the Pelton-Round Butte Hydroelectric Project (Project), constructed at river mile (RM) 100 on the mainstem Deschutes River, creates the primary barrier to anadromous fish attempting to reach spawning and rearing areas in the upper basin. Plans are underway to reinitiate fish passage facilities at the Pelton-Round Butte complex (details in Oregon Steelhead Recovery Plan) and reintroduce steelhead to the upper basin.

*Predation/competition/disease.* This refers to predation, competition, and disease issues in mainstem and estuary that affect all of the Middle Columbia steelhead populations (see Section 6.3.5 of this plan). In addition, the Oregon Steelhead Recovery Plan hypothesizes that the abundance of the Deschutes River Westside population may be limited by competition with a large resident population of rainbow trout.

### **John Day River MPG**

The following are major limiting factors for the John Day River MPG (see also the Oregon Steelhead Recovery Plan [Appendix A]):

*Mainstem passage.* These populations must pass three dams; thus, limiting factors include direct mortality of pre-smolts and smolts at John Day, The Dalles, and Bonneville dams; delayed upstream migration of returning adults; false attraction of returning adults over McNary Dam; and cumulative impact of hydropower system on mainstem and estuary habitat.

*Hatchery related effects.* Hatchery fish straying into natural spawning areas pose risks to genetic traits and productivity of naturally produced steelhead. Concern over competition for resources with wild fish and potential hybridization with natural-origin fish resulted in termination of all hatchery stocking of *O. mykiss* in the John Day River basin in 1997. Most hatchery stray recoveries occur in the lower mainstem John Day River below the North Fork; however, strays have been observed in all populations.

*Tributary habitat.* For all five John Day populations, degraded floodplain and degraded channel structure (key habitat quantity and habitat diversity), altered sediment routing, water quality (temperature), and altered hydrology are limiting factors. For the Lower and Upper Mainstem and South Fork populations, passage obstructions in some of the smaller tributaries are also significant.

*Predation/competition/disease.* This refers to predation, competition, and disease issues in mainstem and estuary that affect all of the Middle Columbia steelhead populations.

### **Umatilla/Walla Walla MPG**

The following are the major limiting factors for the Umatilla/Walla Walla MPG (see also the Oregon Steelhead Recovery Plan [Appendix A] and the Southeast Washington Recovery Plan [Appendix F]):

*Mainstem passage.* The Walla Walla and Touchet populations must pass four major dams; the Umatilla population must pass three.

*Tributary habitat.* For all three populations, water quality (temperature), sediment routing dysfunction, blocked and impaired fish passage, degraded floodplain and channel structure (key habitat quantity and habitat diversity) and hydrologic alterations are limiting factors.

*Hatchery related effects.* The hatchery program on the Umatilla River uses endemic (native) stock and is not currently considered a threat to wild steelhead; however, out-of-DPS strays pose a risk to spawner composition. Non-endemic hatchery fish are considered a potential threat to the Walla Walla wild steelhead population. Currently, data are insufficient to determine whether hatchery effects are a problem for the Touchet River population. An endemic stock program is under development for the Walla Walla and Touchet.

*Predation/competition/disease.* This refers to predation, competition, and disease issues in mainstem and estuary that affect all of the Middle Columbia steelhead populations.

### **Yakima Basin MPG**

The following are primary limiting factors for the Yakima MPG (see also the Yakima Steelhead Recovery Plan [Appendix E]):

*Mainstem passage.* As the farthest upstream populations in the DPS, the Yakima populations must pass four dams and undergo higher exposure to altered habitat and avian and piscine predators in the mainstem Columbia.

*Tributary habitat.* Fish habitat in the Yakima subbasin is substantially influenced by the development of irrigation systems. Limiting factors include altered hydrology (low summer flow because of withdrawals in tributaries and the lower Yakima, scouring peak flows because of degraded watershed conditions, high summer delivery flows in mainstem Yakima and Naches rivers, reduced winter and spring flows due to irrigation storage, delivery, and withdrawals); degraded riparian area and LWD recruitment; blocked and impaired fish passage (primarily due to storage and diversion dams, as well as entrainment in unscreened diversions); altered sediment routing; degraded water quality; loss of historical habitat because of blocked or impaired fish passage; degraded floodplain connectivity and function (loss of off-channel habitat, side channels and connected hyporheic zone); degraded channel structure and complexity; reduced outmigrant survival in the mainstem Yakima.

*Hatchery related effects.* The Yakima populations have the lowest rates of hatchery strays in the DPS, and hatchery effects are not considered a significant limiting factor.

*Predation/competition/disease.* Of the Middle Columbia steelhead populations, the Yakima basin populations have the longest migration through the mainstem Columbia River. They may therefore be more vulnerable to some factors such as avian and piscivorous fish predation. For example, Yakima steelhead, but not the others, are consumed by Caspian tern and double-crested cormorants nesting on islands at the mouth of the Snake River.

### **DPS Recovery Strategy**

NMFS' overall goal for DPS viability, as formulated by the ICTRT and described in Chapter 3 of this plan, is to have all four extant MPGs at viable (low risk) status, with representation of all the major life history strategies present historically, and with the

abundance, productivity, spatial structure and diversity attributes required for long-term persistence.

The ICTRT's current status assessment for the Middle Columbia steelhead DPS and the gaps analysis show that for this DPS, the outlook is optimistic. One population, North Fork John Day, is currently at very low risk or "highly viable." Two populations are currently viable (Deschutes Eastside, Fifteenmile); eleven are at moderate risk, with good prospects for improving. However, the three large populations at high risk (Deschutes Westside, Naches, and Upper Yakima), are important to DPS viability; as a minimum, Deschutes Westside and one of the two large Yakima populations should also reach viable status. These present significant, though not insuperable, challenges. Because of the steelhead's complex life cycle and the many changes that have taken place in its environment, the factors limiting its survival must be addressed in concert, and in an integrated way. The work needs to occur at a regional level, in terms of commitment to actions and funding, and at the local level, population by population.

NMFS' 2006 listing decision called upon Federal, state, and tribal entities to do their best to manage land, hydropower, hatchery, and harvest activities in a manner that would support steelhead recovery. This plan reaffirms those recommendations and adds to them with the contributions of science and consensus building accomplished in the management unit plans.

The recovery strategy for the Middle Columbia steelhead DPS is made up of the following elements:

- Address the limiting factors for each major population group and population, following the recommendations in the 2006 listing decision, making use of the strategies and actions developed in the management unit plans, in concert with the strategies and actions provided in the NMFS 2008 FCRPS Biological Opinion, NMFS Estuary Module, Hatchery and Genetic Management Plans (HGMPs) and *Artificial Production for Pacific Salmon* (Appendix C of Supplemental Comprehensive Analysis, NMFS 2008), fishery management planning through *U.S. v. Oregon* for mainstem fisheries and Fisheries Management Evaluation Plans for tributary fisheries.
- Address and coordinate DPS-wide and basin-wide issues through the Middle Columbia Forum (a bi-state, tri-tribe group convened by NMFS to provide input on the development and implementation of the DPS recovery plan).
- Coordinate research, monitoring, and evaluation throughout the range of the DPS
- Conduct periodic comprehensive reviews of new information generated through the research, monitoring, and evaluation program. Adapt the strategies and actions as appropriate to achieve the recovery plan goals.

NMFS believes that if this strategy is implemented and the biological response is as expected, the Middle Columbia steelhead DPS is likely to achieve viable status within 25 to 50 years.

### **Recovery Strategies for the Four Major Population Groups**

These summaries of recovery strategies for the four major population groups are drawn from the management unit plans and the ICTRT’s status assessment (ICTRT 2008).

#### **Cascades Eastern Slope Tributaries MPG**

<b>Population</b>	<b>ICTRT Risk Status</b>
Fifteenmile Creek (Oregon)	Viable
Deschutes Eastside (Oregon)	Viable
Klickitat (Washington)	(provisional) Moderate risk – insufficient data, hatchery influence
Rock Creek (Washington)	(provisional) High risk – insufficient data
Deschutes Westside (Oregon)	High risk
White Salmon (Washington)	Functionally extirpated
Crooked River (Oregon)	Extirpated

*Primary limiting factors and threats (Section 6.4.1):*

- Degraded tributary habitat
- Mainstem passage
- Hatchery related effects
- Blocked migration to historically accessible habitat
- Predation, competition, disease – in mainstem and estuary; possibly also in Deschutes Westside as competition with resident rainbow trout.

*Gap:* The median survival gap (assuming recent ocean and base hydrosystem conditions and 5 percent risk) for the Eastern Cascades MPG is 0.21 (meaning that a 21 percent increase in average life-cycle survival is required to achieve 5 percent risk in a 100-year time period). The gap ranges from –0.34 (Deschutes Eastside) (no gap) to 0.78 (Deschutes Westside) (needs 78 percent improvement). There was not enough information to estimate gaps for the Klickitat or Rock Creek populations.

*Recovery Scenario:* For the Eastern Cascades Slope Tributaries MPG to be considered viable based on the currently extant populations, the Klickitat, Fifteenmile, and both the Deschutes Eastside and Westside populations should reach viable status, with one highly viable. The Rock Creek population should reach “maintained” status (25 percent or less risk level). MPG viability could be further bolstered if reintroduction of steelhead into the Crooked River succeeds and if the White Salmon population is successfully reintroduced to its historical habitat.

*Key actions proposed (Section 7.3.1):*

- Protect, improve, and increase freshwater habitat for steelhead production.  
Improvements to freshwater habitat should be targeted to address specific limiting

factors in specific areas as described in the Oregon Steelhead Recovery Plan and the Washington Gorge plans.

- Reduce straying of out-of-DPS hatchery fish onto natural spawning grounds within the Deschutes subbasin.
- Restore historical passage to the upper Deschutes subbasin including the Westside tributaries and Crooked River above Pelton Round Butte dam complex and the White Salmon River above Condit Dam.
- Improve survival in mainstem and estuary through actions detailed in NMFS Estuary Module (NMFS 2007) and FCRPS Biological Opinion (NMFS 2008) (as summarized in the Hydro Module [NMFS 2006 and in preparation).
- Improve hatchery management to minimize impacts from hatchery releases on naturally produced steelhead within the Deschutes West and East and Klickitat subbasins
- Fill data gaps for better assessment of Klickitat and Rock Creek steelhead populations.
- Coordinate between scientists, planners, and implementers of recovery actions on both sides of the Columbia River for sequencing of recovery actions and monitoring for adaptive management.

**John Day River MPG**

<b>Population</b>	<b>ICTRT Risk Status</b>
North Fork John Day	Highly viable
Upper Mainstem John Day	Moderate risk
Lower Mainstem John Day	Moderate risk
Middle Fork John Day	Moderate risk
South Fork John Day	Moderate risk

*Main limiting factors and threats (Section 6.4.2):*

- Degraded tributary habitat
- Mainstem passage
- Hatchery related effects
- Predation/competition/disease in mainstem and estuary.

*Gap:* The median survival gap for the John Day MPG is 0.09, ranging from –0.49 (North Fork) (no gap) to 0.34 (South Fork) (needs 34 percent improvement in average survival over the life cycle).

*Recovery Scenario:* For the John Day River MPG to reach viable status, the Lower Mainstem John Day River, North Fork John Day River, and either the Middle Fork John Day River or Upper Mainstem John Day River populations should achieve viable status, with one highly viable.

*Key Actions proposed (7.3.2):*

- Protect and improve freshwater habitat conditions and connectivity for steelhead production. Improvements to freshwater habitat should be targeted to address

specific factors in specific areas as described in the Oregon Steelhead Recovery Plan.

- Improve hatchery management to reduce straying from out-of-DPS hatchery fish onto natural spawning grounds within the John Day subbasin.
- Improve survival in mainstem and estuary through actions detailed in NMFS Estuary Module (NMFS 2007) and FCRPS Biological Opinion (NMFS 2008)

**Yakima River MPG**

<b>Population</b>	<b>ICTRT Risk Status</b>
Upper Yakima River	High Risk
Naches River	High Risk
Satus Creek	Moderate Risk
Toppenish Creek	Moderate Risk

*Main limiting factors and threats (Section 6.4.3):*

- Tributary habitat: Influence of major irrigation system development. Altered hydrology; degraded habitat; loss of habitat; impaired fish passage; reduced outmigrant survival in Yakima mainstem.
- Mainstem passage (four dams)

*Status:* The Yakima MPG is currently rated at High Risk. The two largest populations in the drainage (Naches and Upper Yakima) are rated at High Risk; the Satus Creek and Toppenish Creek populations are rated as Maintained.

*Gap:* The median survival gap (assuming recent ocean and base hydrosystem conditions) for the Yakima MPG is 0.77 (needs 77 percent improvement in average survival over the life cycle), ranging from 0.22 (Satus—tributary only) to 1.15 (Upper Yakima). This is the highest median survival gap of the four MPGs in the Middle Columbia steelhead DPS.

*Recovery Scenario:* For the Yakima River MPG to achieve viable status, two populations should be rated as viable, including at least one of the two classified as Large - the Naches River and the Upper Yakima River. The remaining two populations should, at a minimum, meet the Maintained criteria.

*Key actions proposed (Section 7.3.3):*

- Protect and enhance habitat in key tributary watersheds in the Yakima Basin.
- Restore passage to blocked areas in the Naches and Upper Yakima population areas.
- Alter irrigation delivery and storage operations in the Yakima Basin to improve flow conditions for Middle Columbia steelhead and use managed high flows to maintain floodplain habitat.
- Improve channel and floodplain function and reduce predation through the mainstem Yakima and Naches Rivers.

*Proposed Middle Columbia Steelhead ESA Recovery Plan*  
*September 2008*

- Improve survival in the mainstem Columbia and its estuary through actions detailed in NMFS Estuary Module (NMFS 2007) and FCRPS Biological Opinion (NMFS 2008)

**Umatilla/Walla Walla MPG**

<b>Population</b>	<b>ICTRT Risk Status</b>
Umatilla River	Moderate Risk
Walla Walla River	Moderate Risk
Touchet River	High Risk (provisional because of insufficient data)

*Main limiting factors and threats (Section 6.4.3):*

- Mainstem passage (Touchet and Walla Walla populations pass four major dams: the Umatilla population must pass three)
- Tributary habitat
- Hatchery related effects
- Predation/competition/disease

*Gap:* There was sufficient information available to estimate gaps for only two of the three populations within the Umatilla/Walla Walla MPG. Assuming base hydrosystem and recent ocean conditions, the survival gaps for the Umatilla and Walla Walla populations are 0.09 and 0.34, respectively.

*Recovery Scenario:* For the Umatilla/Walla Walla MPG to be viable, two populations should meet viability criteria, and one should be highly viable. The Umatilla River is the only large population, and therefore needs to be viable. Either the Walla Walla River or Touchet River population also need to be viable.

*Key actions proposed (Section 7.3.4):*

- Coordinate between planners, scientists and those implementing recovery actions in Washington and Oregon for sequencing, monitoring, and adaptive management
- Protect and improve freshwater habitat conditions and access for steelhead production. Improvements to freshwater habitat should be targeted to address specific factors in specific areas as described in the Southeast Washington Plan and the Oregon Steelhead Recovery Plan.
- Improve hatchery management to reduce straying from out-of-DPS hatchery fish onto natural spawning grounds within the Umatilla/Walla Walla subbasins.
- Improve survival in mainstem and estuary through actions detailed in NMFS Estuary Module (NMFS 2007) and FCRPS Biological Opinion (NMFS 2008)

## **DPS-Wide and Basin-Wide Issues**

Problems in migratory corridors for juvenile and adult steelhead in tributaries and the mainstem Columbia River should be addressed to improve survival.

### *Impaired fish passage – mainstem Columbia River*

Passage for juvenile steelhead migrating to the ocean and adult steelhead returning to their natal streams is limited primarily by the four Federal dams on the Lower Columbia River mainstem – Bonneville, John Day, The Dalles, and McNary – which are part of the Federal Columbia River Power System (FCRPS). NMFS recently issued a new biological opinion on the effects of FCRPS operations on salmonids, including Middle Columbia River steelhead, and on the predicted results of current and planned improvements to the system that are intended to improve fish survival (NMFS 2008). These improvements are expected to increase the in-river survival of Middle Columbia River juvenile steelhead by 0.3 percent, 5.1 percent, 8.2 percent, and 10.2 percent, depending on the number of dams they must pass. The survival of steelhead adults through the four dams is thought to be relatively high at the present time (about 98.5 percent per project from Bonneville to McNary), and is expected to be maintained or improved.

The current plan for operation of the FCRPS through 2018 (NMFS 2008) contains the following actions intended to address the needs for survival and recovery of ESA-listed salmon and steelhead:

- Continue adult fish passage operations that have resulted in improved survival.
- Improve juvenile fish passage: install removable spillway weirs or similar surface bypass devices at John Day and McNary dams, an extended tailrace spill wall at The Dalles Dam, and various modifications at Bonneville Dam. Passage for steelhead smolts at each of the four Lower Columbia River mainstem projects must reach 96 percent survival.
- Continue and enhance spill for juvenile fish passage.
- Continue reservoir operations and river flows to benefit spring migrating juveniles.
- Develop dry water year operations to better protect migrating juveniles.

### **Dissenting View of State of Oregon Regarding Mainstem Operations**

At the time this proposed recovery plan was being finalized, August 2008, it was the position of the State of Oregon that additional or alternative actions should be taken in mainstem operations of the FCRPS for ESA-listed salmon and steelhead. Some additional or alternative actions recommended by Oregon, while considered, were not included in NOAA's FCRPS Biological Opinion. At this time, Oregon is a plaintiff in litigation against various federal agencies, including NOAA, challenging the adequacy of the measures contained in the current FCRPS Biological Opinion. NOAA is not in agreement with Oregon regarding the need for or efficacy of Oregon's additional or alternative actions.

*Impaired fish passage – tributaries*

Actions to address fish passage in tributaries include:

- Implement locally developed management unit plans to improve fish passage in tributaries.
- Implement recommendations regarding improved passage and flow management by the U.S. Bureau of Reclamation below all its facilities in the Yakima River and the Umatilla River subbasins, provision of fish passage into significant tributaries, and provision of passage over at least two of its storage dams in the Yakima Basin.<sup>9</sup>
- Implement recommendations regarding improvement of fish passage, screening, and flow management in the Walla Walla River subbasin by the U.S. Army Corps of Engineers, and alteration of the flood operating rule for Mill Creek, or alternatively screening the diversion into Bennington Lake.
- Provide passage into the upper Deschutes River above Round Butte/Pelton complex and into the White Salmon River above Condit Dam.

*Degraded tributary habitat*

Measures to improve tributary habitat are contained in the management unit plans and are summarized above by MPG.

*Hatchery-related effects*

The hatchery programs in the Middle Columbia are managed under the Mitchell Act and the *U.S. v. Oregon* process, involving the fisheries co-managers and regulated by NMFS. NMFS is working with the funding agencies and hatchery operators to update and complete Hatchery and Genetic Management Plans (HGMPs) for every hatchery program in the Middle Columbia region as a means of organizing hatchery review and reform. The HGMPs are the basis for NMFS' biological opinions on hatchery programs under sections 7 and 10 and the 4(d) rule, which relate to incidental and direct take of listed species. The HGMPs describe each hatchery's operations and the actions taken to support recovery and minimize ecological or genetic impacts, such as straying and other forms of competition with naturally produced fish.

Evaluating the factors that influence interactions between hatchery fish and naturally produced fish under varying freshwater conditions and ocean conditions is an important area of future research as well as ESA consultations and NEPA review. This is dealt with in more detail in Appendix C of the 2008 FCRPS Biological Opinion (NMFS 2008).

---

<sup>9</sup> The conservation measures in NOAA's 2006 listing decision specifically identify the need for passage at two or more of the storage dams in the Yakima Basin. The Yakima Steelhead Recovery Plan strongly recommends the provision of passage at the storage dams, but notes that the geographic distribution criteria detailed in the plan do provide for combinations of spawning areas that would meet de-listing and short-term recovery thresholds without provision of access above the storage dams (See Appendix E, Section 4.3.7)

The management unit plans propose various actions to reduce deleterious effects of hatcheries on natural production. For example, The Oregon Steelhead Recovery Plan proposes increased marking of Columbia Basin hatchery steelhead with coded-wire tags, and requiring mass marking of all hatchery steelhead releases with, at a minimum, an adipose fin-clip. Regional consensus has not been reached on these strategies, and the Mid-Columbia Forum will continue to pursue agreement on appropriate site-specific strategies. The Klickitat subbasin plan recommends a targeted monitoring program to determine abundance and productivity of natural spawners, determine the proportion of hatchery and wild spawners in the Klickitat subbasin, and determine the adverse effects of Skamania broodstock on the Klickitat population, if any. Further details are available in each management unit plan.

#### *Predation, Competition and Disease*

Major predation issues in the mainstem Columbia River are addressed in Section 7.4.4. NMFS supports the recommendations in the Yakima Steelhead Plan for research and monitoring to track trends in predator populations, understand their impacts on steelhead, and develop appropriate management techniques to reduce predation. Disease in salmonids is caused by multiple factors and probably cannot be directly addressed by recovery actions except in specific instances of known causal factors. It is more likely that nearly all of the recommended recovery actions that improve spawning, rearing, and passage conditions for steelhead and increase the survival, abundance, and productivity of naturally produced fish will result in decreasing incidence of disease.

#### *Harvest*

Although in general harvest is not considered a major threat for the Middle Columbia steelhead DPS, it is important to ensure that impacts from fisheries do not impede recovery, and to perform monitoring and evaluation to verify impacts and reduce existing uncertainties.

### **Time Required and Cost Estimates**

It is important to consider the unique characteristics and challenges of estimating time and cost for salmon and steelhead recovery, given the complex relationship of these fish to the environment and to human activities on land. NMFS estimates that recovery of the Middle Columbia steelhead DPS, like recovery for most of the ESA-listed Pacific Northwest salmon and steelhead, could take 50 to 100 years, although the optimistic view is that it could be much sooner. The management unit plans (Appendices A through F) contain extensive lists of actions to recover the Middle Columbia steelhead DPS populations. These projects were developed using the most up-to-date assessment of Middle Columbia steelhead recovery needs. The management unit plans focus, for the most part, on actions within the next 5 to 15 years. There are many uncertainties involved in predicting the course of recovery and in estimating total costs. Such uncertainties include biological and ecosystem responses to recovery actions as well as long-term and future funding.

**Cost Estimates**

Cost estimates for recovery projects were provided by the management unit entities where available information was sufficient to do so, using the methods described in each management unit plan. No cost estimates are provided for (1) baseline actions (programs that are already in existence and would occur regardless of this recovery plan), which are listed as Not Applicable (N/A); or (2) actions that need costs to be developed, need unit costs, and/or need project scale estimates. These are listed as To Be Determined. Cost figures will be updated as improved information becomes available.

The total estimated cost of restoring habitat for the Middle Columbia steelhead DPS is approximately \$235 million over the initial 5-year period, and approximately \$970 million over 20 to 50 years for all DPS-wide recovery actions for which sufficient information exists upon which to base an estimate (Table ES-4).

**Table ES-4 Summary of Cost Estimates for Habitat Projects for Middle Columbia Steelhead DPS**

Recovery Plan	First 5 Years (\$M)	Project/Program Total (\$M)
Oregon	\$ 103.5	\$ 512.8
Yakima Steelhead <sup>10</sup>	\$ 91.9	\$ 269.3
SE Washington <sup>11</sup>	\$ 25.5	\$ 76.4
Klickitat <sup>12, 13</sup>	\$ 12.9	\$ 103.6
Rock Creek <sup>14</sup>	\$ 1.0	\$ 1.8
White Salmon Steelhead	N/A	\$ 6.5
<b>DPS Totals</b>	<b>\$ 234.8</b>	<b>\$ 970.4</b>

This estimate includes expenditures by local, tribal, state, and Federal governments, private business, and individuals in implementing both capital projects and non-capital work. Administrative costs are embedded in the total management unit cost estimates in Table ES-4. Preliminary research, monitoring and evaluation costs have, in some cases,

<sup>10</sup> The Yakima steelhead plan estimates costs for the first 6 years, and includes preliminary RME cost estimate of \$300K/year. The 5-year estimate is extrapolated from the 6-year cost data .

<sup>11</sup> The SE Washington plan estimates annual steelhead implementation costs at about \$5 million per year. The 5-year estimate is extrapolated by multiplying the annual amount by five.

<sup>12</sup> The Klickitat plan estimates costs for the first 10 years. Five-year estimate extrapolated by dividing the 10-year amount in half.

<sup>13</sup> The Klickitat plan uses a 50-year period to estimate its total project costs.

<sup>14</sup> The Rock Creek plan estimates cost for first 3 years and 10 years. The 5-year estimate is extrapolated from the 3-year value.

been estimated at the management unit level; however, these costs are not included at this time pending completion of research and monitoring plans and further development of each project.

These cost estimates do not include expenses associated with implementing actions within the lower Columbia River, estuary, or Federal Columbia River Power System (FCRPS), first, because of the basin-wide scope and applicability of these actions to all 13 Columbia Basin salmonid species listed as threatened or endangered, and second, because they are considered "baseline actions" that are required through other processes such as section 7 consultations, FERC licensing agreements, and Habitat Conservation Plans, and these costs would occur regardless of the recovery plans. Cost estimates for estuary actions are included in a module that is incorporated into the Plan by reference, and is available on the NMFS Web site: [www.nwr.noaa.gov/Salmon Recovery Planning/ESA Recovery Plans/Other Documents.cfm](http://www.nwr.noaa.gov/Salmon%20Recovery%20Planning/ESA%20Recovery%20Plans/Other%20Documents.cfm). The estuary recovery costs could be further refined following public comment on the module and on the ESA recovery plan for the three listed lower Columbia River ESUs and one listed lower Columbia River steelhead DPS in 2008 or early 2009. Costs for hatchery actions required through other processes such as consultations, permits, and 4(d) Rule implementation are not part of recovery costs reported here because the programs are already in existence or are undergoing required modifications, and the costs would occur regardless of the recovery plan. There are few estimated costs for recovery actions associated with harvest to report at this time. This is because no actions are currently proposed that go beyond those already being implemented through U.S. v. Oregon and other harvest management forums. In the event that additional harvest actions are implemented through these forums, those costs will be added during the implementation phase of this recovery plan. All cost estimates will be refined and updated over time.

Cost estimates from the draft cost chapters in the individual management plans were developed as consistently as possible, in that they all applied guidance provided by NMFS. However, the approaches vary to some degree given the local and independent nature of the planning groups. Costs developed in the management unit plans were estimated using several basic assumptions (i.e., neither baseline costs nor out-of-basin costs were included in the estimates) and used similar cost calculation methodologies. There are, however, differences in the timeframes for cost estimates, whether administrative costs were included or not, and whether research, monitoring and evaluation costs were calculated. The proposed management unit plans' cost estimates will be refined based on public comment, and final cost estimates will be included in the final DPS recovery plan and management unit plans.

### **Potential Effects of Proposed Recovery Actions**


Chapter 9 in this plan presents an analysis of the potential effects of implementing all the proposed recovery actions – in all the “H” sectors – on the abundance and productivity of Middle Columbia River steelhead. This quantitative analysis provides an opportunity to evaluate the efficacy of proposed recovery strategies in light of current knowledge regarding population functioning, including relationships with habitat conditions. Equally important, the quantitative models used in the assessment provide a framework for

productively targeting evaluation efforts as well as for revisiting key assumptions in the future as more information becomes available (e.g., from monitoring responses to initial implementation or from evaluation efforts targeting key uncertainties). Two models were used: Ecosystem Diagnosis and Treatment (EDT) and the All-H-Analyzer (AHA). Methods of analysis are explained in detail in Chapter 9 of this plan.

The analysis indicates, based on the suites of proposed actions in all the sectors, that all Middle Columbia River steelhead populations for which there are adequate data are expected to achieve 95 percent probability of survival (less than 5 percent risk of extinction within 100 years) for abundance/productivity if the most intensive (major) restoration scenarios are implemented and the projected habitat changes are realized after 25 years of implementation. Under minimum restoration scenarios, three populations (Deschutes Westside, Satus, and Upper Yakima) may not achieve less than 5 percent risk for abundance/productivity. However, the Satus population would meet the recovery criteria identified in the Yakima Steelhead Recovery Plan, and even under poor ocean conditions and minimum restoration actions, the abundance and productivity of the other two populations are expected to increase considerably over the baseline.

Figure ES-6 shows the projected (modeled) abundance and productivity of the 14 populations for which there are adequate data (excluding the Rock Creek, Klickitat, and Touchet populations) after 25 years and major restoration actions. The curve represents the abundance and productivity needed to achieve 95 percent probability of survival for the next 100 years.

Figure ES-6. Predicted viability results for Middle Columbia steelhead populations after 25 years of major restoration efforts.



## **Research, Monitoring, and Evaluation**

An important part of the strategy for achieving recovery is the development of a DPS-wide monitoring plan that will support implementation of the recovery plan and long-term adaptive management in response to changes and trends in the data. Two keys to effective implementation are targeting actions to specific areas and monitoring the results of the actions. To achieve these goals, a scientific technical team made up of local scientists, former ICTRT members, and managers will be necessary. The monitoring plan is discussed in more detail in Chapter 10.

## **Adaptive Management**

Adaptive management in salmon recovery planning is a method of decision making in the face of uncertainty. A plan for monitoring, evaluation, and feedback is incorporated into an overall implementation plan so that the results of actions can become feedback on design and implementation of future actions. Adaptive management works by coupling the decision-making process with collection of performance data and its evaluation. Most importantly, it works by offering an explicit process through which alternative strategies to achieve the same ends can be considered.

Within the Middle Columbia Basin, many different organizations, including Federal, state, tribal, local, and private entities, currently conduct programs and actions that could improve Middle Columbia steelhead survival. Development of Middle Columbia regional coordination will be essential for NMFS' future status reviews of the steelhead DPS. Establishing stable funding and staff to produce annual reports is also important.

Management unit planners are developing detailed research, monitoring, evaluation, and adaptive management plans for each management unit based on the principles and concepts laid out in the NMFS draft guidance document, Adaptive Management for Salmon Recovery: Evaluation Framework and Monitoring Guidance ([http://www.nwr.noaa.gov/Salmon-Recovery-Planning/ESA-Recovery-Plans/upload/Adaptive\\_Mngmnt.pdf](http://www.nwr.noaa.gov/Salmon-Recovery-Planning/ESA-Recovery-Plans/upload/Adaptive_Mngmnt.pdf)) The individual RM&E and adaptive management plans will then be combined into a DPS RM&E and adaptive management plan by the Middle-Columbia Science Team. This will ensure that, taken together, the monitoring and evaluation programs for each management unit, combined with monitoring components of the modules incorporated into the plans, address the needs of the entire DPS. The RM&E and adaptive management plans will be used by the Middle Columbia Forum and others to inform and guide projects and programs during implementation.

## **Setting Priorities**

Priorities for recovery actions should be guided by DPS-, MPG-, and population-level recovery criteria and best available scientific information concerning DPS status, the role of the independent populations in meeting DPS and MPG viability, limiting factors and threats, and likelihood of effectiveness of actions. Protection of existing habitat is essential. Issues of funding and local, state, or national support for implementation will also inevitably come into play.

The management unit plans all address these issues in their implementation sections. For recovery actions in the tributaries, priorities will be settled largely at the local level. However, there should be ongoing technical review and support from DPS-level and management unit

science and technical committees. Coordination and communication in “out-of-subbasin” forums will be necessary for actions in the Columbia mainstem, estuary, and/or ocean.


**Coordination/Governance**

Coordination of actions and information-sharing among fisheries biologists, Tribes, local governments, citizen groups, and state and Federal agencies based in both Oregon and Washington is a key component of recovery for this DPS. Benefits of coordination include:

- Dealing with shared migration areas consistently
- Developing coherent MPG-level strategies where populations are in two states (Cascades Eastern Slope MPG; Umatilla/Walla Walla MPG), or the same population is in both states (Walla Walla population)
- Promoting consistent methods for setting recovery objectives, evaluating strategies, and monitoring progress across populations, MPGs, and the DPS

**Middle Columbia Recovery Forum**

This coordination is under development. The recent creation of the Middle Columbia Recovery Forum (Mid-C Forum), to be convened regularly by NMFS, is intended to facilitate such collaboration between scientists and recovery planners on both sides of the Columbia River. Figure ES-7 gives an overview of the relationships between these entities. Chapter 11 of this plan describes in more detail the proposed roles and responsibilities.



**Figure ES-7. Mid-C Recovery Plan Implementation Organizational Structure**

## **Implementation Funding**

Funding for project implementation is currently available from a variety of sources, but it will be an ongoing challenge. The role of the Forum is to ensure management unit plan implementers are aware of potential sources of funds and to advocate for the funding and implementation of actions that benefit all populations in the DPS. The Forum will not supersede decisions made by the individual management unit boards but may promote funding of their projects and programs if requested. Sources of implementation funding include:

- Congressional appropriations to Federal agencies and to Pacific Coastal Salmon Recovery Fund (PCSRF) (through states and tribes).
- Salmon Recovery Funding Board (SRFB) (Washington).
- Oregon Watershed Enhancement Board (OWEB) (Oregon).
- State appropriations (State agencies).
- Northwest Power and Conservation Council Fish and Wildlife Program (States and tribes).
- Federal / state grants.
- Non-profit organization programs and grants.

## **How NMFS Intends to Use the Plan**

Although recovery plans are not regulatory and their implementation is voluntary, they are important tools that help to do the following:

- Provide context for regulatory decisions.
- Guide decision making by Federal, state, Tribal, and local jurisdictions.
- Provide criteria for status reporting and delisting decisions.
- Organize, prioritize, and sequence recovery actions.
- Organize research, monitoring, and evaluation efforts.

NMFS will encourage Federal agencies and non-Federal jurisdictions to take recovery plans under serious consideration as they make the following sorts of decisions and allocate their resources:

- Actions carried out to meet Federal ESA section 7(a)(1) obligations
- Actions that are subject to ESA sections 4d, 7(a)(2), or 10
- Hatchery and Genetic Management Plans and permit requests
- Harvest plans and permits
- Selection and prioritization of subbasin planning actions
- Development of research, monitoring, and evaluation programs
- Revision of land use and resource management plans
- Other natural resource decisions at the state, Tribal, and local levels

NMFS will emphasize recovery plan information in ESA section 7 (a)(2) consultations, section 10 permit development, and application of the section 4(d) rule by considering:

- The importance of affected populations to listed species viability
- The importance of the action area to affected populations and species viability
- The relation of the action to recovery strategies and management actions

*Proposed Middle Columbia Steelhead ESA Recovery Plan  
September 2008*

- The relation of the action to the research, monitoring, and evaluation plan for the affected species

In implementing these programs, recovery plans will be used as a reference and a source of context, expectations, and goals. NMFS staff will encourage the Federal “action agencies” to describe in their biological assessments how their proposed actions will affect specific populations and limiting factors identified in the recovery plans, and to describe any mitigating measures and voluntary recovery activities in the action area.