

Adult Upstream Survival from Bonneville Dam to Lower Granite Dam

| Snake River Steelhead | | | | | | | Snake River Spring Summer Chinook | | | | | | |
|--------------------------------|-------------------------|-------|--------------------------|-------------------------|-------|----------|-----------------------------------|-------|----------|--------------------------|-------|-------|-------|
| Migrated in River as Juveniles | | | Transported as Juveniles | | | | Migrated in River as Juveniles | | | Transported as Juveniles | | | |
| Survival | 95% Confidence interval | | Survival | 95% Confidence interval | | Survival | 95% Confidence interval | | Survival | 95% Confidence interval | | | |
| | Lower | Upper | | Lower | Upper | | Lower | Upper | | Lower | Upper | | |
| 2002 | 75.7% | 76.7% | 82.6% | 68.3% | 70.2% | 77.4% | 2002 | 84.8% | 82.5% | 86.8% | 75.6% | 73.0% | 78.0% |
| 2003 | 78.8% | 73.9% | 89.7% | 75.4% | 78.8% | 87.5% | 2003 | 82.0% | 79.4% | 84.5% | 75.5% | 73.0% | 77.9% |
| 2004 | 80.1% | 76.6% | 85.6% | 70.6% | 70.8% | 80.0% | 2004 | 83.5% | 81.7% | 85.2% | 76.8% | 72.9% | 80.3% |
| 2005 | 73.8% | 74.4% | 85.5% | 74.6% | 74.6% | 84.2% | 2005 | 83.7% | 80.5% | 86.6% | 80.3% | 76.5% | 83.7% |
| 2006 | 76.6% | 77.5% | 92.4% | 67.2% | 64.9% | 80.9% | 2006 | 74.2% | 68.5% | 79.3% | 66.9% | 62.0% | 71.5% |
| 2007 | 71.4% | 73.7% | 89.6% | 61.7% | 71.0% | 85.2% | 2007 | 79.2% | 72.2% | 85.0% | 75.5% | 71.1% | 79.5% |
| 2008 | 64.3% | 79.4% | 85.7% | 55.4% | 63.0% | 68.3% | 2008 | 71.2% | 68.5% | 73.9% | 71.9% | 68.8% | 74.9% |
| 2009 | 67.0% | 73.8% | 75.9% | 64.6% | 73.0% | 76.8% | 2009 | 71.9% | 68.9% | 74.8% | 75.6% | 70.8% | 80.0% |
| 2010 | 69.3% | 75.5% | 78.0% | 65.1% | 71.8% | 76.0% | 2010 | 69.9% | 66.7% | 73.0% | 72.4% | 70.9% | 73.9% |
| 2011 | 68.7% | 75.9% | 78.3% | 63.3% | 71.3% | 75.8% | 2011 | 67.2% | 65.0% | 69.3% | 66.1% | 63.0% | 69.1% |

| Snake River Fall Chinook | | | | | | |
|--------------------------------|-------------------------|-------|--------------------------|-------------------------|-------|-------|
| Migrated in River as juveniles | | | Transported as Juveniles | | | |
| Survival | 95% Confidence interval | | Survival | 95% Confidence interval | | |
| | Lower | Upper | | Lower | Upper | |
| 2002 | 59.6% | 0.0% | 100.0% | | | |
| 2003 | 81.5% | 0.0% | 100.0% | 82.4% | 56.6% | 96.2% |
| 2004 | 77.6% | 0.0% | 100.0% | 78.5% | 66.5% | 87.7% |
| 2005 | 56.6% | 0.0% | 100.0% | 59.7% | 47.0% | 71.5% |
| 2006 | 45.1% | 0.0% | 100.0% | 47.8% | 26.8% | 69.4% |
| 2007 | 66.8% | 0.0% | 100.0% | 53.6% | 33.9% | 72.5% |
| 2008 | 66.9% | 0.0% | 100.0% | 65.9% | 59.8% | 71.6% |
| 2009 | 59.4% | 0.0% | 100.0% | 58.7% | 51.2% | 65.9% |
| 2010 | 70.0% | 0.0% | 100.0% | 66.7% | 64.6% | 68.8% |
| 2011 | 62.5% | 0.0% | 100.0% | 49.0% | 43.9% | 54.1% |